

- **Originalarbeiten**

- 1. Participants of the European Rating Aggression Group (ERAG) including Schneider, B. Social dysfunction and aggression scale (SDAS-21) in generalized aggression and in aggressive attacks, a validity and reliability study. *Int J Methods Psychiatr Res* 1992; 2: 15-29
- 2. Fritze J, Schneider B, Lanczik M. Vaskuläre Risikofaktoren bei affektiven Psychosen. [Vascular risk factors in affective psychoses]. *Krankenhauspsychiatrie* 1993; 4: 14-18
- 3. Fritze J, Schneider B, Lanczik M. Autoaggressive behaviour and cholesterol. *Neuropsychobiology*. 1992; 26(4):180-1
- 4. Gerbaldo H, Helisch A, Schneider B, Philipp M, Benkert O. Subtypes of negative symptoms: the primary subtypes in schizophrenic and non-schizophrenic patients. *Prog Neuro-psychopharmacol Biol Psychiat* 1994; 18: 311-320
- 5. Fritze J, Beneke M, Lanczik M, Schneider B, Walden J. Carbamazepine as adjunct or alternative to lithium in the prophylaxis of recurrent affective disorders. *Pharmacopsychiatry* 1994; 27: 181-185
- 6. Weber B, Petersen WK, Schneider B, Simminger D, Fritze J, Maurer K. Selbstbeurteilung und Verlauf wahn-indizierender dysfunktionaler Kognitionen bei stationär behandelten depressiven Patienten. *Nervenarzt* 1998; 69: 592-599
- 7. Weber B, Fritze J, Schneider B, Simminger D, Maurer K. Computerized self assessment in psychiatric inpatients. Acceptability, feasibility and influence of computer attitude. *Acta Psychiatr Scand* 1998; 98: 140-145
- 8. Schneider B, Philipp M, Müller MJ, Benkert O. Mortalität bei psychiatrischen Erkrankungen - Ergebnisse einer 5-Jahres-Katamnese. I. Unnatürlicher Tod. *Suizidprophylaxe* 1999; 26: 6-17
- 9. Schneider B, Philipp M, Müller MJ, Benkert O. Mortalität bei psychiatrischen Erkrankungen - Ergebnisse einer 5-Jahres-Katamnese. II. Natürlicher Tod. *Suizidprophylaxe* 1999; 26: 18-26
- 10. Markowitsch HJ, Kessler J, Russ MO, Frölich L, Schneider B, Maurer K. Mnestic block syndrome. *Cortex* 1999; 35: 219-230
- 11. Schneider B, Weber B, Frensch A, Stein J, Fritze J. Vitamin D in schizophrenia, major depression and alcoholism. *J Neural Transm* 2000; 107: 839-842
- 12. Schneider B, Müller JM, Philipp M. Mortality in affective disorders. *J Affect Disord* 2001; 65: 263-274
- 13. Schneider B, Philipp M, Müller MJ. Psychopathological predictors of suicide in patients with major depression during a 5-year follow-up. *Eur Psychiatry* 2001; 16: 283-288

- 14. Weber B, Fritze J, Schneider B, Kühner T, Maurer K. Bias in computerized neuropsychological assessment of depressive disorders caused by computer attitude. *Acta Psychiatr Scand* 2002; 105:126-130
- 15. Schneider B, Weber B, Maurer K, Frölich L. Problematik bei der Beurteilung depressiver Symptome Demenzkranker. *Psychiatr Prax* 2002; 29: 76-82
-
- 16. Weber B, Schneider B, Fritze J, Gille B, Hornung S, Kühner T, Maurer K. Acceptance of computerized compared to paper- and – pencil assessment in psychiatric inpatients. *Comput Hum Behav* 2003; 19: 81-93
- 17. Schneider B, Weigmann H, Hiemke C, Weber B, Fritze J. Reduction of clozapine-induced hypersalivation by pirenzepine is safe. *Pharmacopsychiatry* 2004; 37: 43-45
- 18. Schneider B, Maurer K, Sargk D, Heiskel H, Weber B, Frölich L, Georgi K, Fritze J, Seidler A. Concordance of DSM-IV Axis I and II diagnoses by personal and informant's Interview. *Psychiatry Res* 2004; 127: 121-136
- 19. Schneider B, Schnabel A, Sargk D, Maurer K, Weber B, Wetterling T. Detection of alcohol consumption in suicides. *Eur Arch Psychiatry Clin Neurosci* 2005; 255: 1-5
-
- 20. Schneider B, Schnabel A, Weber B, Frölich L, Maurer K, Wetterling T. Nicotine use in suicides: a case-control study. *Eur Psychiatry* 2005; 20: 129-136
-
- 21. Schneider B, Bartusch B, Schnabel A, Fritze J. Achse I-Störungen als Risikofaktoren für Suizid in Abhängigkeit von Alter und Geschlecht. *Psychiatr Prax* 2005; 32: 185-194
-
- 22. Schneider B, Wetterling T, Sargk D, Schneider F, Schnabel A, Maurer K, Fritze J. Axis I-disorders and personality disorders as risk factors for suicide. *Eur Arch Psychiatry Clin Neurosci* 2006; 256: 17-27
- 23. Schneider B, Georgi K, Weber B, Schnabel A, Ackermann H, Wetterling T. Risikofaktoren für Suizid bei Störungen im Zusammenhang mit psychotropen Substanzen. *Psychiatr Prax* 2006; 33: 81-87
-
- 24. Kõlves K, Värnik A, Schneider B, Fritze J, Allik J. Recent life events and suicide: a case-control study in Tallinn and Frankfurt. *Soc Sci Med* 2006; 62: 2887-2896
-
- 25. Wetterling T, Schneider B, Weber B. Dyslipidämie bei langjähriger neuroleptischer Behandlung [Dyslipidemia and schizophrenia]. *Psychiatr Prax* 2007; 34: 223-229
-
- 26. Wetterling T, Schneider B, Weber B. Blutzuckerwerte bei chronisch schizophrenen Patienten unter antipsychotischer Behandlung [Blood glucose in chronic schizophrenic patients treated with antipsychotics]. *Psychiatr Prax* 2007; 34:76-80
-
- 27. Wetterling T, Weber B, Depfenhart M, Schneider B, Junghanns K. Development of a Rating Scale that allows to predict the Severity of Alcohol Withdrawal Syndrome. *Alcohol alcohol* 2006; 41: 611-615
-

- 28. Schneider B, Kölves K, Blettner M, Wetterling T, Schnabel A, Värnik A. Substance use disorders as risk factors for suicide in an Eastern and a Central European city. *Psychiatry Res.* 2009; 165: 263-272
- 29. Weber B, Schneider B, Hornung S, Wetterling T, Fritze J. Computer attitude –measurable in psychiatric inpatients? *Comput Hum Behav* 2008; 24: 1741-1752
- 30. Giegling I, Rujescu D, Mandelli L, Schneider B, Hartmann AM, Schnabel A, Maurer K, De Ronchi D, Möller HJ, Serretti A. Tachykinin receptor 1 variants associated with aggression in suicidal behavior. *Am J Med Genet B Neuropsychiatr Genet* 2007; 144: 757-761
- 31. Serretti A, Mandelli L, Giegling I, Schneider B, Hartmann AM, Schnabel A, Maurer K, Möller HJ, Rujescu D. HTR2C and HTR1A gene variants in German and Italian suicide attempters and completers. *Am J Med Genet B Neuropsychiatr Genet* 2007; 144: 291-299
- 32. Rujescu D, Giegling I, Mandelli L, Schneider B, Hartmann AM, Schnabel A, Maurer K, Möller HJ, Serretti A. NOS-I and -III gene variants are differentially associated with facets of suicidal behavior and aggression-related traits. *Am J Med Genet B Neuropsychiatr Genet* 2008; 147: 42-48
- 33. Gietl A, Giegling I, Hartmann AM, Schneider B, Schnabel A, Maurer K, Möller HJ, Rujescu D. ABCG1 gene variants in suicidal behavior and aggression-related traits. *Eur Neuropsychopharmacol* 2007; 17: 410-416
- 34. Schneider B, Schnabel A, Wetterling T, Bartusch B, Weber B, Georgi K. How do personality disorders modify suicide risk? *J Personal Disord* 2008; 22: 233-245
- 35. Thalmeier A, Dickmann M, Giegling I, Schneider B, Hartmann AM, Maurer K, Schnabel A, Kauert G, Möller HJ, Rujescu D. Gene expression profiling of post mortem orbitofrontal cortex in violent suicide victims. *Int J Neuropsychopharmacol* 2008; 11: 217-228
- 36. Tonelli LH, Stiller J, Rujescu D, Giegling I, Schneider B, Maurer K, Schnabel A, Möller H-J, Postolache TT. Elevated Cytokine Expression in the Orbitofrontal cortex of victims of suicide. *Acta Psychiatr Scand* 2008; 117: 198-206
- 37. Giegling I, Moreno-De-Luca D, Rujescu D, Schneider B, Hartmann AM, Schnabel A, Maurer K, Möller HJ, Serretti A. Dopa decarboxylase and tyrosine hydroxylase gene variants in suicidal behavior. *Am J Med Genet B Neuropsychiatr Genet.* 2008; 147: 308-315
- 38. Schneider B, Wetterling T, Georgi K, Bartusch B, Schnabel A, Blettner M. Smoking differently modifies suicide risk of affective disorders, substance use disorders, and social factors. *J Affect Disord* 2009; 112: 165-173
- 1. 39. Giegling I, Rujescu D, Mandelli L, Schneider B, Hartmann AM, Schnabel A, Maurer K, De Ronchi D, Möller HJ, Serretti A. Estrogen receptor gene 1 variants are not associated with suicidal behavior. *Psychiatry Res* 2008; 160: 1-7
- 2. 40. Klempan T, Rujescu D, Merette C, Himmelman C, Sequeira A, Canetti L, Fiori I, Schneider B, Bureau A, Turecki G. Profiling Brain Expression of the Spermidine/Spermine

N1-Acetyltransferase 1 (1) Gene in Suicide" has been accepted for publication in Neuropsychiatric Genetics. Am J Med Genet B Neuropsychiatr Genet. 2009; 150: 934-943

41. Schneider B, Georgi K. Zur Frage der Verwendbarkeit des Buss-Durkee Aggressionsfragebogens und der Barratt Impulsiveness Skala im Rahmen der Psychologischen Autopsie. Suizidprophylaxe 2009; 139: 176-180

42. Müller B, Georgi K, Schnabel A, Schneider B. Does sport have a protective effect against suicide? Epidemiologia e Psichiatria Sociale 2009; 18: 331-335

43. Giegling I, Chiesa A, Mandelli L, Gibiino S, Hartmann AM, Möller HJ, Schneider B, Schnabel A, Maurer K, De Ronchi D, Rujescu D, Serretti A. Influence of neuronal cell adhesion molecule (NCAM1) variants on suicidal behaviour and correlated traits. Psychiatry Res 2010; 179: 222-225

44. Schneider B, Georgi K, Fritze J, Grebner K. Finden suizidale Suchtkranke Hilfsangebote im Internet? [Do individuals with substance use disorders find information for crisis intervention and suicide prevention resources on the Internet?]. Nervenarzt 2011; 82: 859-865

45. Schneider B, Baumert J, Schneider A, Marten-Mittag B, Meisinger C, Erazo N, Hammer GP, Ladwig KH. The effect of risky alcohol use and smoking on suicide risk. Findings from the German MONICA/KORA-Augsburg Cohort Study. Soc Psych Psych Epid 2011; 46: 1127-1132

- 46. Schneider B, Grebner K, Schnabel A, Georgi K. Is survivors' emotional response dependent on the consequences of the suicide and the received support? Crisis 2011; 32:186-193

- 47. Schneider B, Grebner K, Schnabel A, Georgi K. Do suicides' characteristics influence survivors' emotions? Suicide Life Threat Behav 2011; 41:117-125

48. Schneider B, Grebner K, Schnabel A, Hampel H, Georgi K, Seidler A. H Impact of employment status and work-related factors on risk of completed suicide. A case-control psychological autopsy study. Psychiatr Res 2011; 190: 265-270

49. Wetterling T, Schneider B. Medikamentenmißbrauch bei älteren psychiatrischen Patienten. [Prescription Drug Abuse in Elderly Psychiatric Patients.] Psychiatr Prax. 2012; 39: 275-279

50. Antypa N, Giegling I, Calati R, Schneider B, Hartmann AM, Friedl M, Konte B, Lia L, De Ronchi D, Serretti A, Rujescu D. MAOA and MAOB polymorphisms and anger-related traits in suicidal participants and controls. Eur Arch Psychiatry Clin Neurosci. 2013; 263: 393-403.

51. Wetterling T, Schneider B. Alkoholintoxikation und Suizidalität [Alcohol intoxication and suicidality]. Psychiatr Prax. 2013; 40: 259-263.

52. Schneider B, Lukaschek K, Baumert J, Meisinger C, Erazo N, Ladwig KH. Living alone, obesity, and smoking increase risk for suicide independently of depressive mood. Findings

from the population-based MONICA/KORA Augsburg Cohort Study. *J Affect Disord* 2014; 152-154; 416-421.

•

53. Baumert J, Schneider B, Lukaschek K, Enemy RT, Meisinger C, Erazo N, Dragano N, Ladwig KH. Adverse conditions at the workplace are associated with increased suicide risk. *J Psychiatr Res* 2014; 57: 90-95.

54. Wolfersdorf M, Schneider B, Schmidtke A. Suizidalität: ein psychiatrischer Notfall, Suizidprävention: eine psychiatrische Verpflichtung. [Suicidal behavior: a psychiatric emergency situation, suicide prevention: a psychiatric obligation]. *Nervenarzt*.2015; 86: 1120-1129

55. Sarlon J, Habich O, Schneider B. Elevated Rest Heart Rate in Psychiatric Patients and Different Effects of Psychotropic Medication. *Pharmacopsychiatry* 2016; 49:18-22

56. Lindner R, Schneider B. Psychotherapie der Suizidalität. [Psychotherapy of suicidality]. *Nervenarzt*. 2016; 87: 488-49

57. Schäfer I, Barnow S, Pawils S, CANSAS Study Group (including Schneider B). Substanzbezogene Störungen als Ursache und als Folge früher Gewalt [Substance use disorders as a cause and consequence of childhood abuse. Basic research, therapy and prevention in the BMBF-funded CANSAS-Network]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2016; 59(1): 35-43

58. Balestri M, Crisafulli C, Donato L, Giegling I, Calati R, Antypa N, Schneider B, Marusic D, Tarozi ME, Marusic D, Paragi M, Hartmann AM, Konte B, Marsano A, Serretti A, Rujescu D. Nine differentially expressed genes from a post mortem study and their association with suicidal status in a sample of suicide completers, attempters and controls. *J Psychiatr Res* 2017; 91: 98-104

59. Schäfer I, Pawils S, Driessen M, Härter M, Hillemacher T, Klein M, Muehlhan M, Ravens-Sieberer U, Schäfer M, Scherbaum N, Schneider B, Thomasius R, Wiedemann K, Wegscheider K, Barnow S. Understanding the role of childhood abuse and neglect as a cause and consequence of substance abuse: the German CANSAS network. *Eur J Psychotraumatol* 2017; 8(1): 1304114

60. Melchior H, Hüsing P, Grundmann J, Lotzin A, Hiller P, Pan Y, Driessen M, Scherbaum N, Schneider B, Hillemacher T, Stolzenburg S, Schomerus G, Schäfer I; Cansas Study Group. Substance Abuse-Related Self-Stigma in Women with Substance Use Disorder and Comorbid Posttraumatic Stress Disorder. *Eur Addict Res* 2019; 25(1): 20-29

61. Schäfer I, Lotzin A, Hiller P, Sehner S, Driessen M, Hillemacher T, Schäfer M, Scherbaum N, Schneider B, Grundmann J. A Multisite Randomized Controlled Trial of Seeking Safety vs. Relapse Prevention Training for Women with Co-Occurring Posttraumatic Stress Disorder and Substance Use Disorders. *EJPT Eur J Psychotraumatol*. 2019; 10(1):1577092

62. Gidzgie P, Grundmann J, Lotzin A, Hiller P, Schneider B, Driessen M, Schaefer M, Scherbaum N, Hillemacher T, Schäfer I. The dissociative subtype of PTSD in women with

substance use disorders: Exploring symptom and exposure profiles. *J Subst Abuse Treat.* 2019; 99: 73-79